



CERTIFICATE OF MAILING

I hereby certify that this paper is being deposited with the U.S. Postal Service as first class mail in an envelope addressed to: Commissioner of Patents, P.O. Box 1450, Alexandria VA 22313-1450, on May 5, 2006.

Signature Theodore J. Leitner Date 5/5/06
Name: Theodore J. Leitner

PATENTS
Attorney Docket No. 7459

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of:
Yi Feng Zheng, *et al.*

Serial No.: 10/736,004

Group Art Unit: 1641

Filed: December 15, 2003

Examiner: Shafiqul Haq

Title: Assays for Entactogens

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

DECLARATION OF INVENTOR PURSUANT TO RULE 37 CFR 1.131


I, Yi Feng Zheng, declare that the following statements are true of my own knowledge, or are believed to be true if stated to be based upon information and belief, and if called upon, I would testify to the truth of these statements.

I am a chemist with Dade Behring Inc., the assignee of the above-referenced U.S. Patent Application, Serial No. 10/736,004, filed December 15, 2003 (hereinafter "Application").

I am one of the co-inventors (hereinafter 'Applicant') named in the Application. I have read the Application and I am familiar with the contents of the Application. Moreover, I have read an Office Action from the U.S. Patent and Trademark Office (USPTO) mailed February 6, 2006 (hereinafter "Office Action"), and I am familiar with the contents of the Office Action.

BEST AVAILABLE COPY

Prior to January 23, 2002, my co-inventors and I prepared the following compound:




Attached hereto as Appendix A are notebook pages 26-33 from my Laboratory notebook #4908 at Dade Behring Inc. The notebook pages show the details of the preparation of the above compound.

All of the dates blacked out in Appendix A were prior to January 23, 2002.


I further declare that all statements made herein are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the patent application Serial No. 10/736,004, or any patent issued thereon.

Dated on this 4th day of May 2006 at Wilmington DE.


Yi Feng Zheng

APPENDIX A

31 Feng Zheng 26



Reagents	MW	Weight	mmols	Sources
S-4908-12	289.74	10 mg	0.0345	page 12
EDAC·HCl	191.71	20 mg	0.1043	
NHS ester	115.07	19 mg	0.165	
DMF	(0.6 mL)			
KLH	5.5 x 10 ⁶	20 mg		
PBS	buffer (pH=8.0, 0.1 M, 6 mL)			

Procedure: To a solution of S-4908-12 (10 mg) in DMF (0.6 mL) was added EDAC·HCl (20 mg) and NHS (19 mg). The reaction mixture was stirred under argon at room temperature for 2.5 hours. The activated hapten was added dropwise via a syringe to a well prepared solution of KLH (20 mg) in Na₂HPO₄-Na₂HPO₄ buffer (6 mL, pH=8.0, 0.1 M). The pH value was changed during the addition. 0.1 NaOH was used to adjust pH value to 8.0. After the complete addition, the reaction was allowed to stir overnight at 4°C (16 hours). The reaction mixture then was ready to dialyze. The 31 dialyzing buffer solution was prepared from

Read and Understood by

[Signature]

3: Jeng Zeng

Dulbecco's phosphate buffer saline (Sigma, Cat No D-1408, Lot No: 48H2339) in a ratio of 400 ml of buffer mixing up with 2600 ml of DE Water.

10 mM Na_2HPO_4 - NaH_2PO_4 buffer (pH=7.0) was prepared from stock solution of 0.2 M NaH_2PO_4 and Na_2HPO_4 .

The KLF-Immunogen (reaction mixture) was dialyzed against Dulbecco's phosphate buffer saline.

3 L, 4°C overnight (~16 hours).

3 L, 4°C 24 hours.

3 L, 4°C 40 hours.

10 mM NaH_2PO_4 - Na_2HPO_4 buffer solution (pH=7.0)

3 L, 4°C 3 hours.

3 L, 4°C 4 hours.

14 ml of Immunogen was obtained denoting as

4908-27

Determination of the concentration of KLF-Immunogen by using BCA Protein Concentration Assay.

1) Working Reagent

Reagent A for BCA[®] protein Assay (Pierce Cat No: 73224 Lot No: 9510206)

Reagent B " " " " (Pierce Cat No: 23224 Lot No: 95062567)

Reagent A (25 ml) + Reagent B (0.5 ml) = 25.5 ml

Working Reagent (green)

Read and Understood by

[Signature]

Date

Y. Teng - 2/20/20

28

- 2) Sample Preparation in NaH_2PO_4 - Na_2HPO_4 buffer (pH=7.0) and Sodium borate solution (0.1M)

0.1 ml of Immunogen in 0.2 ml of 0.1M Sodium borate $\frac{1}{3} \times$

- 3) KLT standard

#1 1.30 $\mu\text{g/ml}$ in 0.1 M Sodium borate } duplicated
#2 0.65 $\mu\text{g/ml}$ in 0.1 M Sodium borate } samples
#3 0.325 $\mu\text{g/ml}$ in 0.1 M Sodium borate

- 4) Samples + Standards + Blank = 2 + 1 + 1 = 4 tubes.
each tube contains 0.1 ml of immunogen or standard or blank dilute with 2 ml of working reagent. Incubate at 40°C for 30 minutes

- 5) UV-vis checked OD at 562 nm

	OD $_{562\text{nm}}$	Concentration
#1	1.002	1.30 $\mu\text{g/ml}$
#2	0.5378	0.65 $\mu\text{g/ml}$
#3	0.26675	0.325 $\mu\text{g/ml}$
Immunogen	0.561	2.12 $\mu\text{g/ml}$

→ see standard curve in page 29

$$y = 0.0326 + 0.74756x \quad R = 0.9995$$

$$0.561 = 0.0326 + 0.74756x$$

$$x = 0.7068$$

$$\text{Immunogen concentration} = 3 \times 0.7068 = 2.12 \mu\text{g/ml}$$


Read and Understood by

[Signature]

Date

(continued from page 28) Yi Teng Zhang

OD =
at
 $\lambda = 562 \text{ nm}$



Determination of Hapten Number for KLH

Immunogen

- Reagents
- 1) 0.1 M Sodium borate
 - 2) 0.1% (w/w) TNBS (2,4,6-trinitrobenzene sulphonic acid)
 - 3) 1 M HCl
 - 4) 10% (w/w) SDS (Sodium dodecyl sulfate) solution

- 1) 0.1 M Sodium borate (100 mL) $\text{pH} \approx 9.32$ $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$
MW = 381.37

$$0.1 \text{ M} \times 0.1 \text{ L} = \frac{x}{381.37} \quad x = 3.8137 \text{ g (100 mL H}_2\text{O)}$$

- 2) 10% SDS 10 mL, $\text{CH}_3(\text{CH}_2)_{11}\text{OSO}_3\text{Na}$ MW = 288.22
1 g of SDS in 9 g (9 mL) of H₂O

- 3) 0.1% TNBS solution \cdot TNBS = MW = 293.2
TNBSO₃Na MW = 383.2

Read and Understood by

Yi Feng 2/2/92

30

$$20 \text{ ng} \times \frac{283.5}{293.2} = 26.1 \text{ ng} \quad (20 \text{ mL H}_2\text{O})$$

4) Sample of K₁H Immagen at a concentration 2.12 mg/mL

— 0.5 mL of K₁H Immagen #908-26

— 0.5 mL of 0.1 M Sodium borate

— 0.5 mL of 0.1% TNBS

— 40°C incubation for 2 hours

— 0.5 mL of 1 M HCl

— 0.5 mL 10% SDS

— UV-vis range 200-480 nm, checked OD at $\lambda = 340 \text{ nm}$

5) K₁H

— 0.5 mL of K₁H solution of 1.12 mg/mL

— 0.5 mL of 0.1 M Sodium borate

— 0.5 mL of 0.1% TNBS

— 40°C incubation for 2 hours

— 0.5 mL of 1 M HCl

— 0.5 mL of 10% SDS

— UV-vis range 200-480 nm, checked OD at $\lambda = 340 \text{ nm}$

6) Blank solution

— 1.0 mL of 0.1 M Sodium borate

— 0.5 mL of 0.1% TNBS

— 40°C incubation for 2 hours

— 0.5 mL of 1 M HCl

— 0.5 mL of 10% SDS

— UV-vis range 200-480 nm, checked baseline correction



7) UV-vis scanned at 200-480 nm, checked at $\lambda = 340 \text{ nm}$

see next page	Sample	OD $\lambda = 340 \text{ nm}$	Concentration
f	K ₁ H	0.9674	1.35 mg/mL
	K ₁ H Immagen	0.3580	

Read and Understood by

31

Si Ferry 2nd day



Read and Understood by _____

[Signature]

31 Feb 2020

32

Hapten Number Calculation.

$$[KLH] = \frac{1.50 \text{ mg/ml} \times 0.2}{5.5 \times 10^6} = 4.73 \times 10^{-8}$$

$$A = \epsilon b c \quad \epsilon_{TNBS} = 1.1 \times 10^4 \text{ (TNBS)} \quad b = 1 \text{ cm}$$

$$[C]_{KLH-TNBS} = \frac{0.9674}{1.1 \times 10^4 \times 1 \text{ cm}} = 8.79 \times 10^{-5}$$

TNBS reacted with KLH

$$\frac{8.79 \times 10^{-5}}{4.73 \times 10^{-8}} = 1858$$

$$\text{Immunogen} \rightarrow \text{KLH Immunogen} = \frac{2.10 \text{ mg/ml} \times 0.2}{5.5 \times 10^6} = 7.71 \times 10^{-8}$$

$$A = \epsilon b c$$

$$[C]_{KLH \text{ Immunogen}} = \frac{0.358}{1.1 \times 10^4} = 3.25 \times 10^{-5}$$

TNBS reacted with NH₂ on Immunogen

$$\frac{3.25 \times 10^{-5}}{7.71 \times 10^{-8}} = 421.5 \approx 422$$


$$\text{Hapten No} = 1858 - 422 = 1436$$

$$\text{Hapten \%} = \frac{1436}{1858} \times 100\% = 77.3\%$$

Read and Understood by

[Signature]




Y. Lengyel



ID No: 4908-26
 Name: KLH Immunogen
 Conc: 2.12 mg/ml
 Buffer: 10 mM Na₂HPO₄-NaH₂PO₄, pH = 7.0
 Hapten No: 1436
 Hapten %: 77%
 Amount: 13 mL

page 33
 notebook: 54908

Y. Lengyel



Read and Understood by

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☒ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.